

Time Evolution of Tunneling in Thermal Medium
– Environment-driven Excited Tunneling –

Sh. Matsumoto and M. Yoshimura

Institute for Cosmic Ray Research, University of Tokyo

Kashiwa-no-ha 5-1-5 Kashiwa, Chiba

277-8582 Japan

ABSTRACT

Time evolution of tunneling phenomena proceeding in thermal medium is studied using a standard model of environment interaction. A semiclassical probability formula for the particle motion in a metastable state of one dimensional system put in thermal medium is combined with the formula of quantum penetration factor through a potential barrier, to derive the tunneling rate in medium. Effect of environment, its influence on time evolution in particular, is clarified in a real-time formalism. A nonlinear resonance effect is shown to enhance the tunneling rate at finite times of order $2/\eta$, with η the friction coefficient. In the linear approximation this effect has relevance to the parametric resonance. This effect enhances the possibility of early termination of the cosmological phase transition much prior to the typical Hubble time.

I Introduction

Tunneling phenomena, when they occur in isolation, are genuinely a quantum effect. But when they occur in some surrounding medium, the important question arises as to whether the tunneling rate is enhanced or suppressed by the environment effect. There are already many works on this subject [1], and most past works deal with a system in equilibrium as a whole. The Euclidean technique such as the bounce solution [2] is often used in this context [3]-[8]. We would like to examine this problem by working in a new formalism of the real-time description of tunneling phenomena [9]-[11]. We find it more illuminating to use a real-time formalism instead of the Euclidean method much employed in the literature. An advantage of the real-time formalism is the possibility of identifying an enhanced tunneling rate at a finite time of dynamical evolution of the tunneling [11].

Our formalism is based on separation of a subsystem from thermal environment, and integrating out the environment degrees of freedom. This method is best suited to a (by itself) non-equilibrium system which is immersed in a larger thermal equilibrium state. Detailed properties of the environment and its interaction form with the subsystem are expected to be insensitive to the behavior of the subsystem in question. We use the standard model of environment consisting of an infinite number of harmonic oscillators [3], [12]. In this picture dissipation seen in the behavior of the subsystem is due to our ignorance of the huge environment degrees of freedom. Although the subsystem itself is an open system having an interaction with environment, the entire system including the environment obeys quantum mechanical laws in the closed system.

We work out consequences of the real-time formalism for a simple subsystem in one dimension. The most interesting result that comes out in this study is the resonance enhanced tunneling [11]. The enhancement occurs at a time scale of order $1/\eta$, with η the friction coefficient. The enhanced tunneling probability goes with η like $\eta^{-1.5}$ according to our numerical analysis in the present work. We offer an interpretation of the resonance enhancement due to an energy flow into the system from environment. Furthermore, the enhanced tunneling rate is related to the phenomenon of the parametric resonance. We call this enhanced tunneling as environment-driven excited tunneling (EET). The popular Euclidean approach cannot deal with the finite time behavior of tunneling, hence misses the resonance enhancement at finite times.

Our method, with a certain extension, should be applicable to field theory models. In future we hope to investigate applications to cosmology, in which one may deal with the electroweak first order phase transition relevant to baryogenesis [13]. An important question here is the time scale of tunneling. If the usual Hubble time scale that gives rise to the potential (or more properly the free energy) change, hence regarded in the conventional picture as the termination of the first order phase transition, is replaced by a shorter tunneling time scale of order $1/\eta$, the conventional picture of the first order phase transition may drastically be changed. For instance, bubbles of the true vacuum may be formed by the resonance enhanced tunneling prior to the potential change. Since this process is stochastic, it may enhance the out-of-equilibrium condition necessary for baryogenesis [14], [15]. The bubble formation via EET may also resurrect the old scenario of GUT phase transition [16] which was once discarded due to the graceful exit problem [17].

The rest of this paper is organized as follows. In Section II the model and the basic tool of analysis is explained. In Section III the semiclassical approximation is introduced and the result of the crucial simplification, the linear approximation of the environment variable, is worked out. This approximation is used while the particle in one dimensional potential is in the metastable region, to determine how the particle is excited in the potential well due to the environment interaction. In Section IV detailed numerical analysis in the metastable region is presented, taking a concrete example of the potential. It is shown that excitation to higher levels proceeds via the parametric resonance effect in the linear approximation. In Section V the semiclassical formula in preceding sections is combined with the quantum mechanical barrier penetration factor to derive the tunneling rate in thermal medium. Numerical analysis is then presented, to quantitatively demonstrate the effect of EET. In the final section application to cosmological baryogenesis is briefly discussed. In Appendix we give the friction coefficient along with relevant formulas in the standard model of particle physics, which may be applicable to discussion of the electroweak baryogenesis in the standard model.

II Model and basic tool

We consider as the first step of our investigation the simplest, yet the most basic problem of this kind, one dimensional system described by a potential $V(q)$. The potential $V(q)$, as illustrated in Fig.1, is assumed to have some local minimum at




Figure 1: A typical potential for tunneling in one dimensional system. Curvature parameters, $\tilde{\omega}_R$ at the bottom of the potential, ω_R at the top of the potential, along with the barrier height V_h and the barrier width, characterize a global structure of the potential.

$q = 0$ with $V(0) = 0$, which is separated at the barrier top, $q = q_B (> 0)$, from a global minimum, which we take in the present work to be at $q = \infty$ such that there is no reflection from a wall at the far right. This system is put in thermal medium which acts as an environment. It has infinitely many, continuously distributed harmonic oscillators given by their coordinates $Q(\omega)$. Its coupling to the tunneling system is given by a Hamiltonian [3], [12],

$$q \int_{\omega_c}^{\infty} d\omega c(\omega) Q(\omega). \quad (1)$$

The frequency dependent coupling strength is $c(\omega)$ and ω_c is some threshold frequency. One may imagine a generalized case in which the subsystem variable q is the order parameter for the first order phase transition in cosmology, the homogeneous Higgs field, and the environment oscillator $Q(\omega)$ is a collection of various forms of matter fields coupled to the Higgs field.

The basic dynamical equation in our problem is then

$$\frac{d^2 q}{dt^2} + \frac{dV}{dq} = - \int_{\omega_c}^{\infty} d\omega c(\omega) Q(\omega), \quad (2)$$

$$\frac{d^2 Q(\omega)}{dt^2} + \omega^2 Q(\omega) = - c(\omega) q. \quad (3)$$

One may eliminate the environment variable $Q(\omega)$ from the second equation above, to get the Langevin equation [18],

$$\frac{d^2 q}{dt^2} + \frac{dV}{dq} + 2 \int_0^t ds \alpha_I(t-s) q(s) = F_Q(t). \quad (4)$$

The kernel function α_I here is given by

$$\alpha_I(t) = - \int_{\omega_c}^{\infty} d\omega r(\omega) \sin(\omega t), \quad (5)$$

with

$$r(\omega) = \frac{c^2(\omega)}{2\omega}. \quad (6)$$

By definition the spectral function $r(\omega)$ is positive definite. The random force term $F_Q(t)$ in eq.(4) is linear in initial values of environment variables, $Q_i(\omega)$ and $P_i(\omega) = \dot{Q}_i(\omega)$ at $t = 0$;

$$F_Q(t) = - \int_{\omega_c}^{\infty} d\omega c(\omega) \left(Q_i(\omega) \cos(\omega t) + P_i(\omega) \frac{\sin(\omega t)}{\omega} \right). \quad (7)$$

By taking the thermal bath of temperature $T = 1/\beta$ given by the density matrix,

$$\begin{aligned} \rho_\beta(Q, Q') &= \left(\frac{\omega}{\pi \coth(\beta\omega/2)} \right)^{1/2} \\ &\cdot \exp \left[- \frac{\omega}{2 \sinh(\beta\omega)} \left((Q^2 + Q'^2) \cosh(\beta\omega) - 2QQ' \right) \right], \end{aligned} \quad (8)$$

for each environment oscillator, the following correlation formula in thermal medium is obtained;

$$\langle \{F_Q(\tau), F_Q(s)\}_+ \rangle_{\text{env}} = \int_{\omega_c}^{\infty} d\omega r(\omega) \cos \omega(\tau - s) \coth\left(\frac{\beta\omega}{2}\right) \equiv \alpha_R(\tau - s). \quad (9)$$

The combination of the two real α 's, $\alpha_R(t) + i\alpha_I(t)$, is the well-known real-time thermal Green's function [19] summed over $Q(\omega)$ -oscillators, being added with the weight $c^2(\omega)$.

An often used simplification is the local, Ohmic approximation taking

$$r(\omega) = \frac{\eta\omega}{\pi}, \quad (10)$$

with $\omega_c = 0$, which amounts to [20]

$$\alpha_I(\tau) = \delta\omega^2 \delta(\tau) + \eta \delta'(\tau). \quad (11)$$

This gives the local version of the Langevin equation,

$$\frac{d^2 q}{dt^2} + \frac{dV}{dq} + \delta\omega^2 q + \eta \frac{dq}{dt} = F_Q. \quad (12)$$

The parameter $\delta\omega^2$ is interpreted as a potential renormalization or a mass renormalization in the field theory analogy, since by changing the bare frequency parameter

to the renormalized ω_R^2 the term $\delta\omega^2 q$ is cancelled by a counter term in the potential. On the other hand, η is the Ohmic friction coefficient. This local approximation breaks down both at early and at late times [21], but it is useful in other time regions.

The value of the coefficient of quadratic term of the potential is shifted by environment interaction. We discuss the relation between the renormalized curvature ω_R ($\tilde{\omega}_R$) at the barrier top (the potential bottom) and the physical curvature ω_B (ω_p) here. These physical curvatures ω_B, ω_p are frequently used in the following discussion. We first illustrate this for the curvature at the barrier top. In the vicinity of the barrier top, the renormalized potential is given by $V(q) \simeq -\omega_R^2(q - q_B)^2/2 + V_h + \text{counter terms}$ ($\omega_R > 0$ by definition). The physical curvature ω_B (the tachyon mass in the field theory analogy, which is defined by a positive value) shifted by the environment is determined as a pole position of the Green's function $\langle 0|T[q(t)q(0)]|0\rangle$ at the barrier top, thus as a solution for the isolated pole of the following equation;

$$\omega_B^2 - \omega_R^2 + \int_{\omega_c}^{\infty} d\omega \frac{2\omega_B^2 r(\omega)}{\omega(\omega_B^2 + \omega^2)} = 0. \quad (13)$$

In many past works, the renormalized curvature ω_R was used as a reference curvature to evaluate the environment effect. Due to the positivity of $r(\omega)$, one has $\omega_B^2 < \omega_R^2$ in general. A consequence of this is that the potential barrier is further suppressed by using ω_B instead of taking the renormalized curvature ω_R as the reference point. It is instructive to give an example. In the Ohmic model of eq.(10) the explicit form of the physical curvature ω_B is given by

$$\omega_B = \sqrt{\omega_R^2 + \frac{\eta^2}{4}} - \frac{\eta}{2}. \quad (14)$$

The relation between the renormalized and the physical curvature at the local minimum of the potential is obtained in the same way, to give in the Ohmic model

$$\omega_p = \sqrt{\tilde{\omega}_R^2 - \frac{\eta^2}{4}}. \quad (15)$$

In subsequent discussion we shall specify which parameter we use.

We now turn to description of the macroscopic system determined by dynamical coordinates $q, Q(\omega)$. Any mixture of quantum pure states, namely a quantum statistical state, is described by the density matrix,

$$\rho = \sum_{n=0}^{\infty} w_n |n\rangle\langle n|, \quad (16)$$

where $|n\rangle$ is a linearly independent state vector for a pure quantum state and $0 \leq w_n \leq 1$. As a special case one may deal with a pure state taking $w_n = 0$ for $n \neq 0$. The density matrix that describes the entire quantum mechanical system obeys the equation of motion;

$$i\hbar \frac{\partial \rho}{\partial t} = [H_{\text{tot}}, \rho], \quad (17)$$

where H_{tot} is the total Hamiltonian of the entire system. For the time being, we explicitly write the Planck constant $\hbar$. In the configuration space this density matrix is given by its matrix elements,

$$\rho(q, q'; Q(\omega), Q'(\omega)) = \langle q, Q(\omega) | \rho | q', Q'(\omega) \rangle. \quad (18)$$

Its Fourier transform with respect to relative coordinates, $q - q', Q(\omega) - Q'(\omega)$, is called the Wigner function and is denoted by f_W ;

$$\begin{aligned} f_W(q, p; Q(\omega), P(\omega)) &= \int_{-\infty}^{\infty} d\xi \prod_{\omega} dX(\omega) \exp[-i\xi p - i \int d\omega X(\omega) P(\omega)] \\ &\times \rho(q + \xi/2, q - \xi/2; Q(\omega) + X(\omega)/2, Q(\omega) - X(\omega)/2). \end{aligned} \quad (19)$$

It is easy to show from the master equation (17) that this quantity obeys ;

$$\begin{aligned} \frac{\partial f_W}{\partial t} &= -p \frac{\partial f_W}{\partial q} + \frac{1}{i\hbar} \left\{ V\left(q + \frac{i\hbar}{2} \frac{\partial}{\partial p}\right) - V\left(q - \frac{i\hbar}{2} \frac{\partial}{\partial p}\right) \right\} f_W \\ &- \int d\omega \left(P(\omega) \frac{\partial f_W}{\partial Q(\omega)} + \omega^2 Q(\omega) \frac{\partial f_W}{\partial P(\omega)} + c(\omega) \left(q \frac{\partial}{\partial P(\omega)} + Q(\omega) \frac{\partial}{\partial p} \right) f_W \right). \end{aligned} \quad (20)$$

Reduction to the subsystem by integrating out environment variables $Q(\omega)$ is an essential part for our physical understanding of tunneling in thermal medium. We define the reduced density matrix of the tunneling system by

$$\rho^{(R)}(q, q') = \int dQ(\omega) dQ'(\omega) \delta(Q(\omega) - Q'(\omega)) \cdot \rho(q, q'; Q(\omega), Q'(\omega)). \quad (21)$$

The fact that there is a delta function $\delta(Q(\omega) - Q'(\omega))$ for the environment variable reflects our understanding that one does not observe the environment. A similar reduction is made to the Wigner function, to get the reduced Wigner function $f_W^{(R)}(q, p)$;

$$f_W^{(R)}(q, p) = \int dP(\omega) dQ(\omega) f_W(q, p; Q(\omega), P(\omega)). \quad (22)$$

The reduction to the subsystem here is in spirit opposite to that common in the effective field theory approach; integration over heavy degrees of freedom is

performed. What is integrated out in the present case is the majority of constituents which is taken invariant under frequent thermalizing interaction.

III Basic idea and semiclassical Wigner function

In considering our problem of tunneling in thermal medium we assume that effects of thermal environment are most important in the time period when the particle in one dimensional system remains in the metastable potential well. Thus, we anticipate that at tunneling the simple quantum mechanical formula may directly be used. We thus separate the problem into two parts; first to treat the potential region left to the barrier top, $q < q_B$, by semiclassically integrating out environment variables and then by projecting the derived reduced density matrix onto the energy eigenstate of the subsystem, which makes it possible to convolute the usual barrier penetration factor used in quantum mechanics. In the following two sections we do this and derive a formula for the tunneling rate in thermal medium.

In the semiclassical $\hbar \rightarrow 0$ limit we have in eq.(20)

$$\frac{1}{i\hbar} \left\{ V \left(q + \frac{i\hbar}{2} \frac{\partial}{\partial p} \right) - V \left(q - \frac{i\hbar}{2} \frac{\partial}{\partial p} \right) \right\} \rightarrow \frac{dV}{dq} \frac{\partial}{\partial p}. \quad (23)$$

The resulting equation for the Wigner function,

$$\begin{aligned} \frac{\partial f_W}{\partial t} = & -p \frac{\partial f_W}{\partial q} + \frac{dV}{dq} \frac{\partial f_W}{\partial p} \\ & - \int d\omega \left(P(\omega) \frac{\partial f_W}{\partial Q(\omega)} + \omega^2 Q(\omega) \frac{\partial f_W}{\partial P(\omega)} + c(\omega) \left(q \frac{\partial}{\partial P(\omega)} + Q(\omega) \frac{\partial}{\partial p} \right) f_W \right). \end{aligned} \quad (24)$$

is identical to the classical Liouville equation familiar in classical statistical mechanics. It has an obvious, formal solution;

$$\begin{aligned} f_W(q, p, Q, P) = & \int dq_i dp_i \int dQ_i dP_i f_W^{(i)}(q_i, p_i, Q_i, P_i) \\ & \cdot \delta(q - q_{cl}) \delta(p - p_{cl}) \delta(Q - Q_{cl}) \delta(P - P_{cl}), \end{aligned} \quad (25)$$

where $q_{cl}, Q_{cl}(\omega)$ etc. are the solution of (2), (3) taken as the classical equation with the specified initial condition written in terms of q_i, p_i, Q_i, P_i .

We consider the circumstance under which the tunneling system is initially in a state uncorrelated to the rest of environment. Thus we take the form of the initial Wigner function of a factorized form,

$$f_W^{(i)}(q_i, p_i, Q_i, P_i) = f_{W,q}^{(i)}(q_i, p_i) \times f_{W,Q}^{(i)}(Q_i, P_i), \quad (26)$$

to get the reduced Wigner function after the environment $Q(\omega), P(\omega)$ integration,

$$f_W^{(R)}(q, p; t) = \int dq_i dp_i f_{W,q}^{(i)}(q_i, p_i) K(q, p, q_i, p_i; t), \quad (27)$$

$$K(q, p, q_i, p_i; t) = \int dQ_i dP_i f_{W,Q}^{(i)}(Q_i, P_i) \delta(q - q_{\text{cl}}) \delta(p - p_{\text{cl}}). \quad (28)$$

The quantity q_{cl} is now a solution of the Langevin equation (4), regarded as a function of initial values; $q_{\text{cl}}(q_i, p_i, Q_i, P_i; t)$. The problem of great interest is how further one can practically simplify the kernel function K introduced here.

In many situations one is interested in the tunneling probability when the environment temperature is low enough. At low temperatures of $T \ll$ a typical frequency or curvature scale ω_s of the potential V , one has

$$\omega_s \sqrt{Q_i^2(\omega)}, \sqrt{P_i^2(\omega)} = O[\sqrt{T}] \ll \sqrt{\omega_s}. \quad (29)$$

In the electroweak and the GUT phase transition this corresponds to $T \ll m_H$ (Higgs mass). Expansion of q_{cl} in terms of $Q_i(\omega), P_i(\omega)$ and its truncation to linear terms is then justified. Thus, we use

$$\begin{aligned} \delta(q - q_{\text{cl}}) &= \int \frac{d\lambda_q}{2\pi} \exp[i\lambda_q(q - q_{\text{cl}})] \\ &\approx \int \frac{d\lambda_q}{2\pi} \exp\left[i\lambda_q\left(q - q_{\text{cl}}^{(0)} - \int d\omega \left\{ Q_i(\omega) q_{\text{cl}}^{(Q)}(\omega) + P_i(\omega) q_{\text{cl}}^{(P)}(\omega) \right\}\right)\right], \end{aligned} \quad (30)$$

valid to the first order of $Q_i(\omega), P_i(\omega)$. The zero-th order term is $q_{\text{cl}}^{(0)}$ obeying

$$\frac{d^2 q_{\text{cl}}^{(0)}}{dt^2} + \left(\frac{dV}{dq}\right)_{q_{\text{cl}}^{(0)}} + 2 \int_0^t ds \alpha_I(t-s) q_{\text{cl}}^{(0)}(s) = 0, \quad (31)$$

while the coefficient functions of the first order terms are $q_{\text{cl}}^{(Q)}(\omega), q_{\text{cl}}^{(P)}(\omega)$ obeying equations later shown, eq.(37). A similar expansion for $\delta(p - p_{\text{cl}})$ using $p_{\text{cl}}^{(0)}, p_{\text{cl}}^{(Q)}(\omega), p_{\text{cl}}^{(P)}(\omega)$, also holds.

An alternative justification of the expansion (30) is to keep $O[c(\omega)]$ terms consistently in the exponent since both $q_{\text{cl}}^{(Q)}(\omega)$ and $q_{\text{cl}}^{(P)}(\omega)$ are of this order. Thus, both in the low temperature and in the weak coupling case our subsequent analysis follows.

With (30) one can proceed further. Gaussian integral for the variables $Q_i(\omega), P_i(\omega)$, first, and then for λ_q, λ_p can be done explicitly with eq.(30), when one takes the thermal, hence Gaussian density matrix for the initial environment variable:

$$f_{W,Q}^{(i)}(Q_i, P_i) = \frac{1}{\pi} \tanh \frac{\beta\omega}{2} \exp\left[-\tanh \frac{\beta\omega}{2} \left(\omega Q_i^2 + \frac{P_i^2}{\omega}\right)\right], \quad (32)$$

The result of this Gaussian integral leads to an integral transform [10] of the Wigner function, $f_W^{(i)}$ (initial) $\rightarrow f_W^{(R)}$ (final), using the following form of the kernel function;

$$K(q, p, q_i, p_i; t) = \frac{\sqrt{\det \mathcal{J}}}{2\pi} \exp \left[-\frac{1}{2} (q - q_{\text{cl}}^{(0)}, p - p_{\text{cl}}^{(0)}) \mathcal{J} \begin{pmatrix} q - q_{\text{cl}}^{(0)} \\ p - p_{\text{cl}}^{(0)} \end{pmatrix} \right], \quad (33)$$

where the matrix elements of $\mathcal{J}_{ij}^{-1} = I_{ij}$ are given by

$$I_{11}(t) = \frac{1}{2} \int_{\omega_c}^{\infty} d\omega \coth \frac{\beta\omega}{2} \frac{1}{\omega} |z(\omega, t)|^2, \quad (34)$$

$$I_{22}(t) = \frac{1}{2} \int_{\omega_c}^{\infty} d\omega \coth \frac{\beta\omega}{2} \frac{1}{\omega} |\dot{z}(\omega, t)|^2, \quad (35)$$

$$I_{12}(t) = \frac{1}{2} \int_{\omega_c}^{\infty} d\omega \coth \frac{\beta\omega}{2} \frac{1}{\omega} \Re(z(\omega, t) \dot{z}^*(\omega, t)) = \frac{\dot{I}_{11}}{2}, \quad (36)$$

where $z(\omega, t) = q_{\text{cl}}^{(Q)}(\omega, t) + i\omega q_{\text{cl}}^{(P)}(\omega, t)$, and $\dot{z}(\omega, t) = p_{\text{cl}}^{(Q)}(\omega, t) + i\omega p_{\text{cl}}^{(P)}(\omega, t)$.

Quantities that appear in the integral transform are determined by solving differential equations; the homogeneous Langevin equation for $q_{\text{cl}}^{(0)}$, eq.(31) and an inhomogeneous linear equation for $z(\omega, t)$ and $\dot{z}(\omega, t)$,

$$\frac{d^2 z(\omega, t)}{dt^2} + \left(\frac{d^2 V}{dq^2} \right)_{q_{\text{cl}}^{(0)}} z(\omega, t) + 2 \int_0^t ds \alpha_I(t-s) z(\omega, s) = -c(\omega) e^{i\omega t}. \quad (37)$$

A similar equation as for $z(\omega, t)$ holds for $\dot{z}(\omega, t)$. The initial condition is

$$q_{\text{cl}}^{(0)}(t=0) = q_i, \quad p_{\text{cl}}^{(0)}(t=0) = p_i, \quad z(\omega, t=0) = 0, \quad \dot{z}(\omega, t=0) = 0. \quad (38)$$

The physical picture underlying the formula for the integral transform, eq.(27) along with (33), should be evident; the probability at a phase space point (q, p) is dominated by the semiclassical trajectory $q_{\text{cl}}^{(0)}$ (environment effect of dissipation being included in its determination by the third term of eq.(31)) reaching (q, p) from an initial point (q_i, p_i) whose contribution is weighed by the probability $f_W^{(i)}$ initially given. The contributing trajectory is broadened by the environment interaction with the width factor $\sqrt{I_{ij}}$. The quantity I_{11} given by (34), for instance, is equal to $\overline{(q - q_{\text{cl}}^{(0)})^2}$; an environment driven fluctuation under the stochastic force $F_Q(t)$.

We note that in the exactly solvable model of inverted harmonic oscillator potential the identical form of the integral transform eq.(33) was derived [9] without resort to the semiclassical approximation. Explicit form of the classical and the fluctuation functions ($q_{\text{cl}}^{(0)}$ and $z(\omega, t)$) is given there, which is valid beyond the Ohmic approximation.

It would be instructive to write solutions in the harmonic approximation taking the frequency at the local potential minimum with $\omega_p = \sqrt{\tilde{\omega}_R^2 - \frac{\eta^2}{4}}$, eq.(15), the potential is given by $V(q) = \frac{1}{2} \omega_p^2 q^2$. Then the zero-th order term $q_{\text{cl}}^{(0)}(t)$ and $z(\omega, t)$ are obtained analytically, and given by

$$q_{\text{cl}}^{(0)} = u(t) q_i + v(t) p_i, \quad (39)$$

$$u(t) = (\cos \omega_p t + \frac{\eta}{2\omega_p} \sin \omega_p t) e^{-\eta t/2}, \quad v(t) = \frac{\sin \omega_p t}{\omega_p} e^{-\eta t/2}, \quad (40)$$

$$z(\omega, t) = \frac{c(\omega)}{\omega^2 - \tilde{\omega}_R^2 - i\omega\eta} \left(e^{i\omega t} - u(t) - i\omega v(t) \right). \quad (41)$$

We used the local Ohmic approximation such as

$$2 \int_0^t ds \alpha_I(t-s) z(\omega, s) \rightarrow \eta \dot{z}(\omega, t). \quad (42)$$

The function $z(\omega, t)$ has a Breit-Wigner form for a small η . In particular, the infinite time ($t \rightarrow \infty$) limit of the fluctuation has

$$I_{11}(\infty) = \int_{\omega_c}^{\infty} d\omega \coth \frac{\beta\omega}{2} \frac{r(\omega)}{(\omega^2 - \tilde{\omega}_R^2)^2 + \eta^2 \omega^2}. \quad (43)$$

The friction η appears as the resonance width in this formula. The real part of the pole position in the frequency integral such as (43), $\omega_p \equiv \Re$ (pole position), is $\omega_p = \sqrt{\tilde{\omega}_R^2 - \frac{\eta^2}{4}}$.

The harmonic approximation is excellent at late times, but at early and intermediate times nonlinear resonance effects are important as discussed in [11]. The nonlinear effect will be taken into account in discussion of the finite time behavior of the tunneling rate.

A convenient form of the fluctuation formula is derived by separating the zero-temperature result along with $\coth \beta\omega/2 = 1 + 2/(e^{\beta\omega} - 1)$. We denote the temperature dependent part of the fluctuation by $I_{ij}^{(T)}$. The zero-temperature part $I_{ij}^{(0)}$ must be renormalized. After performing renormalization the zero-temperature part is approximated by result of the harmonic approximation. The renormalized fluctuation is then well approximated using the physical curvature, ω_p , eq.(15). We shall use the same notation I_{ij} for the renormalized fluctuation in the rest of our paper.

IV Numerical analysis in the metastable region; relevance of parametric resonance




Figure 2: The tunneling potential used in numerical calculation.

We point out a relation of the semiclassical Wigner function in the preceding section to the parametric resonance, as briefly mentioned in [11]. Presence of anharmonic terms is inevitable in any realistic tunneling potential. This introduces a non-trivial oscillating term in the coefficient function $\left(\frac{d^2V}{dq^2}\right)_{q_{\text{cl}}^{(0)}}$ of the fluctuation equation, (37), thus giving a differential equation akin to the Mathieu equation. Depending on whether or not the relevant parameters fall into the instability band of this Mathieu-type equation, the fluctuation may increase indefinitely in the zero friction limit [22]. What happens is more subtle; the parameters fall right on the boundary of stability and instability bands. We shall show this by taking an example of the tunneling potential.

The potential we take (illustrated in Fig.2) is given by

$$\begin{aligned} V(x) &= \frac{\tilde{\omega}_R^2}{2} x^2 + g \tilde{\omega}_R^3 x^4, \quad (x < x_b) \\ &= V_h, \quad (x_b < x < x_f), \quad = -V(\infty), \quad (x_f < x). \end{aligned} \quad (44)$$

The two parameters V_h and x_b are related since we took a particular quartic potential in the metastable well.

The fluctuation $z(\omega, t)$ in the Ohmic approximation follows the equation;

$$\ddot{z}(\omega, t) + V''(q_{\text{cl}}^{(0)}) z(\omega, t) + \eta \dot{z}(\omega, t) = -c(\omega) e^{i\omega t}. \quad (45)$$

The potential derivative here is

$$V''(q_{\text{cl}}^{(0)}) = \tilde{\omega}_R^2 + 12g \tilde{\omega}_R^3 (q_{\text{cl}}^{(0)})^2. \quad (46)$$

Taking the zero friction limit $\eta = 0$ gives the following related homogeneous equation,

$$\ddot{z}(\omega, t) + [h + 2\theta (q_{\text{cl}}^{(0)})^2] z(\omega, t) = 0, \quad (47)$$




Figure 3: Band structure for relevant fluctuation. Shaded areas correspond to the instability band. 1st band is enlarged separately (the left figure), where the crossed point corresponds to our model relation, eq.(48).

where

$$h = \tilde{\omega}_R^2, \quad \theta = 6g\tilde{\omega}_R^3. \quad (48)$$

The equation for $q_{\text{cl}}^{(0)}$ is as follows:

$$\ddot{q}_{\text{cl}}^{(0)} + V'(q_{\text{cl}}^{(0)}) = 0 \quad (49)$$

This gives a periodic solution for $q_{\text{cl}}^{(0)}$. A notable feature of the homogeneous equation in the $\eta = 0$ limit (47) is that $q_{\text{cl}}^{(0)}$ appearing in the coefficient function is periodic, hence it gives rise to solutions either of the Bloch wave type or of the parametric resonance type [22]. Despite of the definite relation, eq.(48), we regard (h, θ) as free parameters to thoroughly investigate the stable-unstable band structures, because that would more definitely clarify relation between the band structure and the relevant parameter region of this model.

For simplicity we analyzed the problem assuming

$$g = 0.01, \quad V_h = 5\omega_p, \quad \omega_p = \sqrt{\tilde{\omega}_R^2 - \eta^2/4}. \quad (50)$$

In a relevant region of

$$0 < \theta < 2\tilde{\omega}_R^3, \quad 0 < h < 5\tilde{\omega}_R^2, \quad (51)$$

which includes the model region (48), we numerically checked the time evolution of the homogeneous solution of eq.(47). We observed the power-law increase of the fluctuation. Indefinite increase of the fluctuation $z(\omega, t)$ as time increase implies that the assumed parameter lies either within the unstable band or on the boundary between the bands. The linear rise $z(\omega, t) \propto t$, as indicated by our numerical analysis




Figure 4: Time evolution of the function $y(t)$, a solution of eq.(54). The amplitude of the function $y(t)$ with the friction $\eta = 0.005\omega_p$ is depicted.

suggests that the parameter is on the boundary. The result of the band structure is given in Fig.3, and the relevant parameter eq.(48) (corresponding to the crossed point) is just on the boundary between stable and 1st unstable bands. This is consistent with that the linear rise, as is expected in the case of the exact Mathieu function on the boundary [23].

For this numerical calculation we had to choose an initial condition for $q_{\text{cl}}^{(0)}$. We took several conditions for investigating band structures, and we always obtained the same result of the linear power-law increase.

We next consider effects of the friction η . The inclusion of the friction gives a modification of eq.(47) to

$$\ddot{z}_h(t) + [h + 2\theta (q_{\text{cl}}^{(0)})^2]z_h(t) + \eta\dot{z}_h(t) = 0, \quad (52)$$

The effect of the friction term $\eta\dot{z}_h(t)$ is studied by changing the variable $z_h(t)$ to $y(t)$,

$$y(t) = z_h(t)e^{\eta t/2}. \quad (53)$$

Then the equation for $y(t)$ becomes

$$\ddot{y} + [(h - \eta^2/4) + 2\theta (q_{\text{cl}}^{(0)})^2]y = 0. \quad (54)$$

On the other hand, the equation for $q_{\text{cl}}^{(0)}$ is

$$\ddot{q}_{\text{cl}}^{(0)} + V'(q_{\text{cl}}^{(0)}) + \eta\dot{q}_{\text{cl}}(0) = 0. \quad (55)$$

When the effect of friction for the behavior of the zero-th order $q_{\text{cl}}^{(0)}$ is small, the equation for y is nearly of the parametric resonance type. In this case then, the




Figure 5: Examples of ω dependence of $z(\omega, t)$. The quantity $\omega_p |z(\omega, t)|^2 + |\dot{z}(\omega, t)|^2 / \omega_p$ both a) at early and b) late times are shown. The unit of time is ω_p^{-1} .

behavior of $z(\omega, t)$ is rather simple; it is a product of the linearly rising function ($y(t) \sim t$) and the exponentially decreasing function ($e^{-\eta t/2}$). The rate of the exponential decrease is $\eta/2$, hence this product function has a maximum at a time around $2/\eta$. This seems essentially what we observe in numerical computation for $z(\omega, t)$.

With the presence of the friction the function y approaches the behavior of simple harmonic oscillator as time increases since $q_{cl}^{(0)} \rightarrow 0$ by friction. This means, as seen in Fig.4, that the linear increase of y is saturated at the time of order $2/\eta$. The net effect of the friction is simply the minor change of a coefficient of A/η , where A is close to 1.7.

The effect of the presence of the inhomogeneous term in eq.(45) is as follows. This term acts as an external force, and gives a resonance effect when the frequency ω is close to the inherent frequency of the system. We illustrate this point. In Fig.5 ω -dependence of $\omega_p |z(\omega, t)|^2 + |\dot{z}(\omega, t)|^2 / \omega_p$ at a particular time is shown, and one observes, besides the simple resonance, higher modes due to the nonlinear effect. Without the strong resonance effect like this one, the fluctuation $z(\omega, t)$ is very small, since we have the initial condition, $z(\omega, 0) = 0 = \dot{z}(\omega, 0)$. Thus, the environment acts here as a force of resonant kick from the zero amplitude, which then makes possible the parametric resonance effect to work, typically very important for large amplitude oscillation. In Fig.5a), the growth of 1st ($\omega \sim \omega_p$) and 2nd ($\omega \sim 3\omega_p$) mode is shown. Other higher modes (3rd, 4th, $\dots$) are also amplified in the same manner as in the 2nd mode case at $t \sim 360\omega_p^{-1}$. On the other hand, these higher modes disappear at late times, and only the first mode remains as seen in Fig.5b).




Figure 6: Time evolution of a) the fluctuation $\omega_p I_{11}^{(T)} + I_{22}^{(T)}/\omega_p$ and b) the subsystem energy $\langle H_{sys.} \rangle - (T = 0 \text{ part})$. A temperature $T = 0.5\omega_p$ is taken with four cases of the friction η (in the unit of ω_p).

Thus, the growth of higher modes is a finite time effect.

We now discuss the behavior of the frequency integrated fluctuation $I(t)$ in eqs.(34)-(36). The increase of the fluctuation due to the parametric resonance gives an interesting time evolution of I . In Fig.6a), the time evolution of the quantity $\omega_p I_{11}^{(T)} + I_{22}^{(T)}/\omega_p$ is shown for a few values of the friction η . This function depends on the initial value, q_i, p_i , for which we took a typical value in the ground state at the potential minimum.

What is the physical picture behind the resonance enhancement? We would like to present a suggestive interpretation. We first derive a relation between the fluctuation here and the subsystem energy defined by

$$\langle H_{sys.} \rangle \equiv \left\langle \frac{p_{cl}^2}{2} + \frac{\omega_p^2}{2} q_{cl}^2 \right\rangle. \quad (56)$$

By making the approximation on q_{cl} and p_{cl} , namely taking up to linear terms of the initial environment value as used in eq.(30), this equation is reduced to

$$\begin{aligned} \langle H_{sys.} \rangle &= (T = 0 \text{ part}) \\ &+ \frac{1}{2} \int_0^\infty \frac{dp_i dq_i}{2\pi} f_W^{(i)}(q_i, p_i) \left\{ I_{22}^{(T)}(q_i, p_i; t) + \omega_p^2 I_{11}^{(T)}(q_i, p_i; t) \right\}. \end{aligned} \quad (57)$$

Thus, the temperature dependent part of the subsystem energy is given by the fluctuation averaged over initial values. In short, the subsystem energy is driven by the fluctuation. If initial values for $\langle q_{cl}^2(0) \rangle$, and $\langle p_{cl}^2(0) \rangle$ are small (in the $\omega_p = 1$ unit), the zero-temperature part ($T = 0$) part is close to $0.5\omega_p$ (zero-point fluctuation

of the system). In Fig.6b) the quantity H_{sys} – (T = 0 part) is shown, taking as the initial state the ground state of the harmonic oscillator at the potential bottom. The ground state we use is characterized by

$$f_W^{(i)}(q_i, p_i) = \frac{1}{\pi} \exp \left[-\omega_p q_i^2 - \frac{p_i^2}{\omega_p} \right]. \quad (58)$$

The time at which the maximum of H_{sys} occurs is $1.7/\eta$, which is the same as for that of the fluctuation, thus indicating a close correlation between the fluctuation and our subsystem energy $\langle H_{sys.} \rangle$. We shall have more to say on an important role of $\langle H_{sys.} \rangle$ towards the end of the next section.

Implication to the tunneling rate seems obvious; the parametric resonance we observe here enhances excitation to higher energy states at the potential region left to the barrier top in the quantum mechanical terminology. This would then enhance the tunneling rate from the metastable state of the potential well. We shall demonstrate this in the following section.

V Tunneling rate formula

Energy eigenstates are special among pure quantum states, since they time-evolve only with phase factors. Thus, if one takes for the projection $|n\rangle\langle n|$ in eq.(16) the energy eigenstate $|n\rangle$ of the total Hamiltonian, the density matrix does not change with time. On the other hand, if one takes the energy eigenstate of the subsystem Hamiltonian, then the density matrix changes solely due to the environment interaction. It is thus best to use a pure eigenstate of, or its superposition of, the subsystem Hamiltonian, when one wishes to determine how the tunneling rate is modified in thermal medium.

We thus perform projection onto the energy eigenstate of the tunneling system. The projected probability onto energy eigenstate of the subsystem Hamiltonian is in general given by

$$\mathcal{W}_E(t) = \langle E | \rho^{(R)} | E \rangle = \int dq dq' dp \varphi_E(q) \varphi_E^*(q') e^{ip(q'-q)} f_W^{(R)}\left(\frac{q+q'}{2}, p; t\right), \quad (59)$$

where $\varphi_E(q)$ is the wave function of eigen energy E .

Our basic idea is to limit application of this formula to the left region of the metastable state and to use the semiclassical formula for the Wigner function

$f_W(q, p; t)$ derived in the preceding section. Explicit p integration in (59) gives

$$\begin{aligned} \mathcal{W}_E(t) = & \int dq dq' \frac{dq_i dp_i}{2\pi} \varphi_E(q) \varphi_E^*(q') f_W^{(i)}(q_i, p_i) \frac{1}{\sqrt{2\pi I_{11}}} \\ & \exp \left[-\frac{1}{2I_{11}} \left(\frac{q+q'}{2} - q_{\text{cl}}^{(0)}(t) \right)^2 - \frac{\det I}{2I_{11}} (q-q')^2 \right. \\ & \left. + i(q' - q) \left(p_{\text{cl}}^{(0)}(t) + \frac{I_{12}}{I_{11}} \left(\frac{q+q'}{2} - q_{\text{cl}}^{(0)}(t) \right) \right) \right]. \end{aligned} \quad (60)$$

The tunneling probability in medium is then obtained by multiplying the well-known formula of the barrier penetration factor.

Unless the energy eigenstate near the barrier top is important for computation of the tunneling rate, one may approximate the wave function in the left region to $q = q_B$ by that of the harmonic oscillator at the potential bottom, $q = 0$. The wave function $\varphi_E(q)$ is thus approximated by using the Hermite polynomial H_n as

$$\varphi_E(q) = \left(\frac{\sqrt{\omega_p}}{\sqrt{\pi} 2^n n!} \right)^{1/2} H_n(\sqrt{\omega_p} q) e^{-\omega_p q^2/2}, \quad (61)$$

where ω_p is the shifted frequency at the local potential minimum, eq.(15).

To compute the projected probability $\mathcal{W}_n(t)$, we introduce the generating function;

$$\mathcal{W}(t, z) = \sum_{n=0}^{\infty} \mathcal{W}_n(t) z^n. \quad (62)$$

The following formula of the Hermite polynomial, the Mehler's formula [24],

$$\sum_{n=0}^{\infty} \frac{(z/2)^n}{n!} H_n(x) H_n(y) = (1-z^2)^{-1/2} \exp \left[\frac{2xyz - (x^2 + y^2)z^2}{1-z^2} \right]. \quad (63)$$

is useful. From eqs.(60), (61), (63), one has Gaussian integrals for q and q' , resulting in

$$\mathcal{W}(t, z) = \int \frac{dq_i dp_i}{2\pi} f_W^{(i)}(q_i, p_i) \sqrt{\frac{1}{\det \tilde{A}}} \exp \left[-\frac{1-z}{2 \det \tilde{A}} (q_{\text{cl}}^{(0)}, p_{\text{cl}}^{(0)}) \tilde{A} \begin{pmatrix} q_{\text{cl}}^{(0)} \\ p_{\text{cl}}^{(0)} \end{pmatrix} \right], \quad (64)$$

$$\tilde{A} = \begin{pmatrix} (1-z)(I_{22} - \frac{\omega_p}{2}) + \omega_p & -(1-z)I_{12} \\ -(1-z)I_{12} & (1-z)(I_{11} - \frac{1}{2\omega_p}) + \frac{1}{\omega_p} \end{pmatrix}. \quad (65)$$

Quantities I_{ij} are to be computed at finite time t . The important constraint of the probability conservation $\sum_{n=0}^{\infty} \mathcal{W}_n(t) = 1$ is automatically satisfied due to




Figure 7: Time evolution of the projected probability for the ground, the second excited, and the fourth excited state. The case of $T = 0.5\omega_p$ and $\eta = 0.005\omega_p$ is depicted. For comparison the result in the harmonic approximation is shown.

$\mathcal{W}(t, 1) = 1$. For precise computation, one expands this generating function to powers of z , and identifies the projected probability $\mathcal{W}_n(t)$ by (62).

We next clarify the relation,

$$\sum_{n=0}^{\infty} E_n \mathcal{W}_n(t) = \langle H_{sys.} \rangle \quad (66)$$

where $\langle H_{sys.} \rangle$ is calculated by (57). This relation was numerically checked to be correct, as might be expected. We have no analytic proof of this relation, although there is no doubt on the correctness of this relation.

All numerical calculations were performed by using the concrete example of the tunneling potential eq.(44) throughout this section. We assume the subsystem is in the ground state at the initial moment, and use eq.(58) as the initial Wigner function. Some results were compared with that of the Harmonic approximation discussed in the last of the section III. The effect of the non-linear term is clarified by this comparison, because the term is neglected in the Harmonic approximation.

In Fig.7, a few examples of the projected probability are shown, assuming the initial ground state; $\mathcal{W}_n(0) = \delta_{n0}$. It is clearly observed that excitation to higher energy levels is correlated to evacuation of the ground state. The infinite time limit of $\mathcal{W}_n(\infty)$ coincides with the result of the harmonic approximation at the potential bottom,

$$\mathcal{W}_n(\infty) = (1 - e^{-\beta\omega})e^{-n\beta\omega} + \mathcal{O}(\eta), \quad (67)$$




Figure 8: a) Time evolution of the tunneling rate $\Gamma(t)$. A temperature $T = 0.5\omega_p$ is taken with four cases of the friction η . b) Ratio of the maximal tunneling rate at $t \sim \mathcal{O}(2/\eta)$ to the $\Gamma(\infty)$ (a traditional one) is depicted as a contour map in the $\eta - T$ plane.

as is expected. We note that the subsystem thermalizes at the infinite time despite that it was initially in the ground state of a harmonic oscillator, hence a non-equilibrium state.

The tunneling rate $\Gamma(t)$ per unit time is roughly the tunneling probability $\times \frac{\omega_p}{2\pi}$, since $\omega_p/2\pi = 1/\text{period}$ for the periodic motion in the left potential well. Thus,

$$\Gamma(t) = \frac{\omega_p}{2\pi} \sum_{n=0}^{\infty} \mathcal{W}_n(t) |T(E_n)|^2, \quad (68)$$

$$|T(E)|^2 \approx \exp \left[-2 \int_{q_1(E)}^{q_2(E)} dx \sqrt{2(\tilde{V}(x) - E)} \right] \theta(V_h - E) + \theta(E - V_h). \quad (69)$$

In computing the tunneling rate we use a modified potential $\tilde{V}$ which is obtained from V by replacing the curvature parameter ω_R by $\omega_B = \sqrt{\omega_R^2 + \eta^2/4} - \eta/2$. The reason for this is that one should compare the tunneling rate with and without the environment effect by using effectively the same values of parameters for the two cases. This modification explains the bulk of the environment effect, the environment suppression factor proposed by [3], as shown in [9]. In equations above $q_i(E)$ are turning points separating the subbarrier region.

In Fig.8a) the tunneling rate $\Gamma(t)$ is shown for a few values of the friction. The tunneling rate is observed to become maximal at the time $t \sim 1.7/\eta$. A salient feature, and the most important result of the present work, is that the tunneling rate is enhanced around the time of order $2/\eta$, which is caused by non-linear resonance effects. The enhanced tunneling rate at $t \approx \mathcal{O}(2/\eta)$ is clearly related to the power-

law rise of the fluctuations I_{ii} , as observed in our previous paper [11] and as already explained. The maximal enhancement factor for the tunneling rate is plotted in Fig.8b), relative to the harmonic approximation. A large enhancement is obtained for small η and small T values. In the harmonic approximation an almost monotonic behavior of time evolution is evidently observed, thus the rate being given by $\Gamma(\infty)$. We may summarize what happens, in the following way. The non-linear resonance inherent to the tunneling potential excites higher modes that have larger tunneling probabilities. This effect is however terminated by the friction caused by environment interaction, which is the origin of the final decrease of the tunneling rate at large times.

We finally study at a quantitative level correlation between the tunneling rate $\Gamma(t)$ and the tunneling system energy $H_{sys}(t)$. In order to evaluate the correlation quantitatively, we define the correlation coefficient r used in statistics;

$$r = \frac{\langle \Gamma H_{sys} \rangle - \langle \Gamma \rangle \langle H_{sys} \rangle}{\langle \Gamma \rangle \langle H_{sys} \rangle}, \quad \langle A \rangle \equiv \frac{1}{\text{Time}} \int_0^{\text{Time}} dt A(t). \quad (70)$$

The correlation coefficient r was numerically calculated and we obtained results that can be summarized by $r = 0.99 \sim 1$ in the parameter region of temperature and friction, $0.2 \leq T/\omega_p \leq 1.0$ and $0.001 \leq \eta/\omega_p \leq 0.02$. This proves the correlation between the two quantities.

We next turn to the discussion of the survival probability, the probability that the metastable state remains in the same initial state. This quantity $P(t)$ is defined by

$$\dot{P}(t) = -\Gamma(t)P(t), \quad P(0) = 1. \quad (71)$$

This gives

$$P(t) = \exp \left[- \int_0^t dt' \Gamma(t') \right]. \quad (72)$$

In Fig.9a) the survival probability is depicted along with the result of the harmonic approximation of the potential bottom. The effect of EET is unquestionable.

A traditional definition of the lifetime t_{trad} which many authors use is

$$t_{\text{trad}} = \Gamma^{-1}(\infty), \quad (73)$$

while in the case of time dependent tunneling one may effectively use t_{LT} given by the e-holding time,

$$P(t_{\text{LT}}) = e^{-1}. \quad (74)$$




Figure 9: a) Time evolution of the survival probability $P(t)$. The case of $T = 0.5\omega_p$ and $\eta = 0.005\omega_p$ is depicted. For comparison the result in the harmonic approximation is shown. b) Ratio of the lifetime as defined in eq.(74) to the traditional one is depicted as a contour map in the $\eta - T$ plane.

In Fig.9b) a contour plot of $t_{LT}/t_{\text{trad.}}$ is shown in the parameter (η, T) space. For small values of η and T a substantial reduction of the lifetime t_{LT} is observed, which means the early termination of the tunneling.

We hope that we have convinced the reader of the power and the usefulness of EET, although our idea is vindicated by detailed numerical analysis which is necessarily limited to special cases.

VI Discussion

There may be many applications of environment-driven excited tunneling (EET) presented in this paper. We shall only mention two possible applications to cosmology, an enhancement mechanism of electroweak baryogenesis and a possible resurrection of the old inflationary scenario of the GUT phase transition. Furthermore in the present discussion we shall focus on one aspect of the baryogenesis condition; the out-of-equilibrium condition [14], [15]. Needless to say, there are many problems to be solved in actual application of these ideas to realistic models.

Electroweak baryogenesis is expected to occur if the electroweak phase transition is of a strong first order. The first order phase transition usually proceeds via the bubble formation. Inside the bubble the true electroweak broken phase is realized in which the baryon number is effectively conserved. On the other hand, outside the

bubble the universe remains in the high temperature symmetric phase in which the baryon number is not conserved via sphaleron related processes[25]. This mismatch inside and outside the bubble gives rise to a possibility of electroweak baryogenesis, although details of a promising mechanism are yet to be worked out. Thus, the strong first order of the phase transition has been considered to be a requisite for the out-of equilibrium condition necessary for the baryogenesis.

In the ordinary circumstance, namely in the standard model, the strong first order phase transition requires a small Higgs mass, which is inconsistent with observation. Along with the small CP violation effect of Kobayashi-Maskawa theory, this excludes [26] the possibility [27] of the standard model as a viable model for baryogenesis. Thus, a more complicated model such as two Higgs doublet model needs to be invoked for the electroweak baryogenesis. Regardless of whether an extended Higgs model is needed, we would like to point out that the environment-driven excited tunneling (EET) found in this paper has a chance of enhancing the out-of equilibrium condition necessary for the baryogenesis.

Although not much discussed in the literature, there is an important factor to consider for the out-of equilibrium condition. This is the factor Γt_H , where Γ is the nucleation rate of the bubble and $t_H = 1/H$ is the Hubble time. In the usual scenario of the first order phase transition the effectiveness of electroweak baryogenesis is in proportion to this quantity. The nucleation rate Γ in the usual estimate without EET is exponentially suppressed by the potential barrier, and the factor Γt_H is much less than 1, which means that the first order phase transition is never completed by a merger of nucleated bubbles. Under this circumstance it is expected that the phase transition is terminated by proceeding to disappearance of the local minimum of the symmetric phase so that the condition for the first order phase transition is not met. The effectiveness of the out-of equilibrium condition is then reduced by a factor proportional to the quantity Γt_H .

On the other hand, with the aid of EET, the factor Γt may become of order unity at some finite time, and the phase transition may be completed much prior to the Hubble time, as discussed in previous sections. One should check in comparison to the Hubble time the new time scale $2/\eta$ which is roughly the time at the maximum for the tunneling rate. The factor Γt thus may become of order unity at $t \sim 2/\eta$. In this case there is no reduction factor like Γt_H .

We shall check numerical relation between two time scales; t_H and $2/\eta$. The size of the friction η in the standard model is of the order of the Higgs mass m_H times

some power of coupling factors, while the Hubble rate is given by

$$H = 1/t_H = \sqrt{\frac{4\pi^2}{45}} N \frac{T^2}{m_{pl}}, \quad N = 106.75. \quad (75)$$

Calculation, as summarized in Appendix, indicates that

$$\eta \simeq \frac{y_t^2}{32\pi} \omega_p \tanh\left(\frac{\omega_p}{4T}\right), \quad \omega_p^2 = V''(0) \simeq 0.2 \times (T^2 - m_H^2 - 0.04v^2), \quad (76)$$

where y_t is the top Yukawa coupling, and $V(\phi)$ is the effective potential of the standard model at finite temperature, with v the vacuum expectation value of the Higgs field. Thus, the Hubble rate is much smaller than η by the inverse Planck factor since $\frac{m_H}{m_{pl}} \ll y_t^2$.

One may expect that even in the case of weak first order phase transition the quantum tunneling enhanced by EET may give rise to a sufficient condition for the out-of-equilibrium in the early phase of the phase transition. Much remains to be seen with this regard.

Another interesting effect of EET concerns resurrection of the old inflationary scenario of the GUT phase transition [16]. The old inflationary scenario was once rejected because the bubble formation proceeds too slowly, hence leaving a very inhomogeneous universe [17]. The essential reason of this problem, called graceful exit problem, is that the exponentially suppressed tunneling rate is too small such that bubbles of true vacuum do not merge sufficiently to give rise to a homogeneous universe which the inflationary scenario intends to derive. Our mechanism of enhanced tunneling may help this situation. The problem must however be addressed quantitatively; one has to obtain an early termination of the first order phase transition to promote merger of bubbles, yet has to obtain an sufficient inflation to solve intended problems. We leave this quantitative study to future.

Acknowledgments

This work is supported in part by the Grant-in-Aid for Science Research, Ministry of Education, Science and Culture, Japan (No.13640255 and No.13135207)

Appendix

A-1 Effective potential and critical temperature

We shall first give the finite temperature effective potential of the standard model at 1-loop order, assuming that the Higgs mass is much smaller than other particle masses of the standard model such as the top quark, the weak boson, thus $m_H \ll m_t, m_Z, m_W$. This is taken as a necessary condition for the first order phase transition. This condition however is not met in reality. Nevertheless, we take this case as an important illustration for more complicated cases.

The zero temperature part of the effective potential [28] is as follows;

$$\begin{aligned} V_0(\phi) &= \frac{m_H^2}{8v^2}(\phi^2 - v^2)^2 + 2Bv^2\phi^2 - \frac{3}{2}B\phi^4 + B\phi^4 \log\left(\frac{\phi^2}{v^2}\right), \\ B &= \frac{3}{64\pi^2 v^4} (2m_W^4 + m_z^2 - 4m_t^2) \simeq -0.005, \end{aligned} \quad (77)$$

where $v = 246\text{GeV}$ is the vacuum expectation value (VEV) of the Higgs field, and m_a is the mass of a particle species a . The finite temperature effect is then given by the free energy [29],

$$\begin{aligned} V_T(\phi) &= \frac{T^4}{2\pi^2} \left[6I_B\left(\frac{m_W\phi}{vT}\right) + 3I_B\left(\frac{m_z\phi}{vT}\right) - 6I_F\left(\frac{m_t\phi}{vT}\right) \right], \\ I_{B,F}(a^2) &= \int_0^\infty dr r^2 \log \left[1 \mp \exp\left(-\sqrt{r^2 + a^2}\right) \right]. \end{aligned} \quad (78)$$

The Higgs contribution has been neglected, due to the assumption of $m_H \ll$ masses of other particles.

According to [29], we may make the high temperature expansion of the effective potential with $T \ll v$, to get

$$V(\phi) = V_0(\phi) + V_T(\phi) \simeq D(T^2 - T_0^2)\phi^2 - ET\phi^3 + \frac{\lambda(T)}{4}\phi^4, \quad (79)$$

where the dimensionless coefficients are given by

$$\begin{aligned} D &= \frac{2m_W^2 + m_z^2 + 2m_t^2}{8v^2} \simeq 0.2, \\ E &= \frac{2m_W^3 + m_z^3}{4\pi v^3} \simeq 0.01, \\ T_0^2 &= \frac{m_H^2 - 8Bv^2}{4D} \simeq 1.25(m_H^2 + (50\text{GeV})^2), \\ \lambda(T) &= \frac{m_H^2}{2v^2} - \frac{3}{16\pi^2 v^4} \left(2m_W^4 \log \frac{m_W^2}{a_B T^2} + m_z^4 \log \frac{m_z^2}{a_B T^2} - 4m_t^4 \log \frac{m_t^2}{a_F T^2} \right). \end{aligned} \quad (80)$$

Here $a_B = 2 \log 4\pi - 2\gamma \simeq 3.91$, $a_F = 2 \log \pi - 2\gamma \simeq 1.14$. The critical temperature is computed from these, to give

$$T_c^2 = \frac{T_0^2}{1 - E^2/(\lambda(T)D)} \simeq \frac{T_0^2}{1 - (8GeV/m_H)^2}. \quad (81)$$

At the value of T_c two local minima coincide, while T_0 indicates where the symmetric phase of $\varphi = 0$ disappears as the local minimum. It is thus important to note that the condition $T_c > T_0$ for the first order phase transition is met.

A-2 Friction coefficient in the standard model

The friction coefficient η is computed using the method of [30],[31]. According to [31], it is possible to identify the two-body bilinear operator of standard model fields as the environment variable $Q(\omega)$. The most dominant contribution to the friction η comes from the top loop diagram where the top-Yukawa coupling is of the form, $y_t \phi \bar{t}t/\sqrt{2}$. Thus, the Langevin equation of motion for the zero-mode Higgs field ϕ_0 becomes

$$\ddot{\phi}_0 + V'(\phi_0) + 2 \int_0^t ds \Im \alpha(t-s) \phi_0(s) = F(t), \quad (82)$$

$$\langle \{F(t), F(s)\} \rangle_{\text{env}} = \Re \alpha(t-s), \quad (83)$$

$$\alpha(x_0) = \frac{y_t^2}{2} \int d^3x \langle \bar{t}t(x) \bar{t}t(0) \rangle_{\text{env}} = \int d\omega \frac{r_H(\omega)}{e^{\beta\omega} - 1} e^{i\omega x_0}. \quad (84)$$

The real-time Green's function $\langle \bar{t}t(x) \bar{t}t(0) \rangle_{\text{env}}$ can be obtained by the analytic continuation of the imaginary-time Green's function [19], and it is easy to calculate the imaginary-time Green's function by using the Matsubara formalism. The necessary spectral weight $r_H(\omega)$ is given by

$$r_H(\omega) = \frac{y_t^2}{2} \int \frac{d^3p}{(2\pi)^3} \frac{\omega^2}{4p^2} \delta(\omega - 2p) (1 - 2n_p), \quad (85)$$

where $p = |\vec{p}|$ is the momentum magnitude and $n_p = (e^{\beta p} + 1)^{-1}$, assuming that the top quark is massless in the symmetric phase around $T \approx T_0$.

In the vicinity of the false vacuum, namely $\varphi \approx 0$, the behavior of the zero-mode Higgs field exhibits the exponentially damped oscillation. The solution of the homogeneous zero-mode Higgs field equation (82) is approximately obtained by

$$\begin{aligned} \phi_0(t) &\simeq 2 \int_0^\infty d\omega \frac{r_H(\omega) \sin(\omega t)}{(\omega^2 - \omega_p^2) + \pi^2 r_H^2(\omega)} \\ &\simeq 2 \int_0^\infty d\omega \frac{r_H(\omega_p) \sin(\omega t)}{(\omega^2 - \omega_p^2) + \pi^2 r_H^2(\omega_p)} = \sin(\omega_p t) \exp\left(-\frac{\pi r_H(\omega_p)}{2\omega_p} t\right), \end{aligned} \quad (86)$$

where ω_p is the curvature of the false vacuum ($\omega_p^2 = V''(0)$), and the friction coefficient in this case is computed from eq.(85). The result is

$$\eta = \frac{\pi}{\omega_p} r(\omega_p) = \frac{y_t^2}{32\pi} \omega_p \tanh \frac{\beta \omega_p}{4} . \quad (87)$$

References

- [1] For a review, P. Hanggi, P. Talkner, and M. Borkovec, *Rev. Mod. Phys* **62**, 251(1990).
- [2] S. Coleman, *Phys. Rev.* **D15**, 2929(1977); C.G. Callan and S. Coleman, *Phys. Rev.* **D16**, 1762(1977); J.S. Langer, *Ann. Phys. (N.Y.)* **41**, 108(1967); M.B. Voloshin, I.Yu. Kobzarev, and L.B. Okun, *Sov. J. Nucl. Phys.* **20**, 644(1975).
- [3] A.O. Caldeira and A.J. Leggett, *Ann. Phys. (N.Y.)* **149**, 374(1983).
- [4] I.K. Affleck, *Phys. Rev. Lett.* **46**, 388(1981).
- [5] A.I. Larkin, and Yu.N. Ovchinnikov, *Sov. Phys. -JETP* **59**, 420(1984).
- [6] H. Grabert, P. Olschowski, and U. Weiss, *Phys. Rev.* **B36**, 1931(1987) and references therein.
- [7] D. Waxman and A.J. Leggett, *Phys. Rev.* **B32**, 4450(1985).
- [8] W. Zwegler, *Z. Phys.* **B51**, 301(1983).
- [9] Sh. Matsumoto and M. Yoshimura, "Dynamics of barrier penetration in thermal medium: exact result for inverted harmonic oscillator", hep-ph/0006019, and *Phys. Rev.* **A63**, 012104(2000).
- [10] Sh. Matsumoto and M. Yoshimura, "Quantum tunneling in thermal medium", hep-ph/0008025.
- [11] Sh. Matsumoto and M. Yoshimura, "Resonance Enhanced Tunneling", hep-ph/0009230, and *Phys. Lett.* **B495**, 223(2000).
- [12] R.P. Feynman and F.L. Vernon, *Ann. Phys. (N.Y.)* **24**, 118(1963).

- [13] For a review, A.G. Cohen, D.B. Kaplan, and A.E. Nelson, *Annu. Rev. Nucl. Part. Sci.* **43**, 27(1993).
- [14] A.D. Sakharov, *Pis'ma Zh. Eksp. Teor. Fiz.* **5**, 32(1967) [*JETP Lett.* **5**, 24(1967)].
- [15] M. Yoshimura, *Phys. Rev. Lett.* **41**, 281(1978); **42**, 746(E)(1979); *Phys. Lett.* **B88**, 294(1979).
- [16] A. Guth, *Phys. Rev.* **D23**, 347(1981); K. Sato, *Mon. Not. R. Astron. Soc.* **195**, 467(1981).
- [17] A. Guth and E. Weinberg, *Nucl. Phys.* **B212**, 321(1983).
- [18] G.W. Ford, J.T. Lewis, and R.F. O'Connell, *Phys. Rev.* **A37**, 4419(1988).
- [19] A.L. Fetter, and J. D. Walecka, *Quantum theory of many-particle systems*, (McGraw-Hill, New York 1971).
- [20] A.O. Caldeira and A.J. Leggett, *Physica A* **121**, 587(1983).
- [21] V. Ambegaokar, Ber. Bunsenges, *Phys. Chem.* **95**, 400(1991); I. Joichi, Sh. Matsumoto, and M. Yoshimura, *Phys. Rev.* **A57**, 798(1998), and references therein.
- [22] L. Landau and E. Lifschitz, *Mechanics* (Pergamon, Oxford, 1960), p. 80.
- [23] N. W. McLachlan, *Theory and application of Mathieu functions*, (Oxford University Press), p. 141.
- [24] Batemann Manuscript Project II, *Higher transcendental functions* (R.E. Krieger Pub. Co., Malabar, Fla, 1981), p. 194.
- [25] V.A. Kuzmin, V.A. Rubakov and M.E. Shaposhnikov, *Phys. Lett.* **B155**, 36(1985); **B191**, 171(1987).
- [26] M.B. Gavela, M. Lozano, J. Orloff, O. Pene, and C. Quimbay, *Nucl.Phys.* **B430**, 345(1994); **B430**, 382(1994); P. Huet, and E. Sather, *Phys. Rev.* **D51**, 379(1995).
- [27] G. R. Farrar, and M.E. Shaposhnikov, *Phys. Rev. Lett.* **70**, 2833(1993); Erratum-ibid.**71**, 210(1993); *Phys. Rev.* **D50**, 774(1994).

- [28] S. Coleman and E. Weinberg, *Phys. Rev.* **D7**, 1888(1973).
- [29] M. Dine, R.G. Leigh, P. Huet, A. Linde, and D. Linde, *Phys. Rev.* **D46**, 550(1992).
- [30] H.A. Weldon *Phys. Rev.* **D28**, 2007(1983).
- [31] I. Joichi, S. Matsumoto, and M. Yoshimura, *Prog. Theo. Phys* **98**, 9(1997).